

Therapist:	
Phone:	

# **ARTHROSCOPIC SLAP LESION REPAIR (TYPE II)**

## I. <u>Phase I – Immediate Postoperative Phase</u> "Restrictive Motion" (Day 1 to Week 6)

<u>Goals</u>: Protect the anatomic repair Prevent negative effects of immobilization Promote dynamic stability Diminish pain and inflammation

#### Week 0-2:

- Sling for 4 weeks
- Sleep in immobilizer for 4 weeks
- Elbow/hand ROM
- Hand gripping exercises
- Passive and gentle active assistive ROM exercise
  - Flexion to 60 degrees (Week 2: Flexion to 75 degrees)
  - Elevation in scapular plane to 60 degrees
  - ER/IR with arm in scapular plane
  - ER to 10-15 degrees
  - IR to 45 degrees

\*\*NO active ER or Extension or Abduction

- Submaximal isometrics for shoulder musculature
- NO isolated Biceps Contractions
- Cryotherapy, modalities as indicated

#### Week 3-4:

- Discontinue use of sling at 4 weeks
- Sleep in immobilizer until Week 4
- Continue gentle ROM exercises (PROM and AAROM)
  - Flexion to 90 degrees
  - Abduction to 75-85 degrees
  - ER in scapular plane and 35° abd to 25-30 degrees
  - IR in scapular plane and 35° abd to 55-60 degrees
  - \*\*NOTE: Rate of progression based on evaluation of the patient.

AOKC.NET

- No active ER, Extension or Elevation
- Initiate rhythmic stabilization drills
- Initiate proprioception training
- Tubing ER/IR at 0 degrees Abduction
- Continue isometrics
- Continue use of cryotherapy





## Week 5-6:

- Gradually improve ROM
  - Flexion to 145 degrees
  - ER at 45 degrees abduction: 45-50 degrees
  - ER at 45 degrees abduction: 55-60 degrees
  - At 6 weeks begin light and gradual ER at 90° abduction progress to 30-40° ER
- May initiate stretching exercises
- May initiate light (easy) ROM at 90 degrees Abduction
- Continue tubing ER/IR (arm at side)
- PNF manual resistance
- Initiate Active Shoulder Abduction (without resistance)
- Initiate "Full Can" Exercise (Weight of Arm)
- Initiate Prone Rowing, Prone Horizontal Abduction
- NO Biceps Strengthening

## II. Phase II – Intermediate Phase: Moderate Protection Phase (Week 7-14)

<u>Goals</u>: Gradually restore full ROM (week 10) Preserve the integrity of the surgical repair Restore muscular strength and balance

#### Week 7-9:

- Gradually progress ROM:
  - Flexion to 180 degrees
  - ER at 90 degrees abduction: 90-95 degrees
  - IR at 90 degrees abduction: 70-75 degrees
- Continue to progress isotonic strengthening program
- Continue PNF strengthening
- Initiate Throwers Ten Program

## Week 10-12:

- May initiate slightly more aggressive strengthening
- Progress ER to Throwers Motion
  - ER at 90 degrees abduction: 110-115 in throwers (Week 10-12)
- Progress isotonic strengthening exercises
- Continue all stretching exercises
  \*\*Progress ROM to functional demands (i.e. overhead athlete)
- Continue all strengthening exercises

## III. Phase III – Minimal Protection Phase (Week 14-20)

Goals: Establish and maintain full ROM

## AOKC.NET



Improve muscular strength, power and endurance Gradually initiate functional activities

#### Criteria to enter Phase III:

- 1) Full non-painful ROM
- 2) Satisfactory stability
- 3) Muscular strength (good grade or better)
- 4) No pain or tenderness

#### Week 14-16:

- Continue all stretching exercises (capsular stretches)
- Maintain Throwers Motion (Especially ER)
- Continue strengthening exercises:
  - Throwers Ten Program or Fundamental Exercises
  - PNF Manual Resistance
  - Endurance training
  - Initiate light plyometric program
  - Restricted sport activities (light swimming, half golf swings)

#### Week 16-20:

- Continue all exercise listed above
- Continue all stretching
- Continue Throwers Ten Program
- Continue Plyometric Program
- Initiate interval sport program (throwing, etc)
  \*\*See interval Throwing Program

## IV. Phase IV – Advanced Strengthening Phase (Week 20-26)

<u>Goals</u>: Enhanced muscular strength, power and endurance Progress functional activities Maintain shoulder mobility

## Criteria to enter Phase IV:

- 1) Full non-painful ROM
- 2) Satisfactory static stability
- 3) Muscular strength 75-80% of contralateral side
- 4) No pain or tenderness

#### Week 20-26:

- Continue flexibility exercises
- Continue isotonic strengthening program
- PNF manual resistance patterns

## AOKC.NET



- Plyometric strengthening
- Progress interval sport programs

## V. Phase V – Return to Activity Phase (Month 6 to 9)

<u>Goals</u>: Gradual return to sport activities Maintain strength, mobility and stability

#### Criteria to enter Phase V:

- 1) Full functional ROM
- 2) Muscular performance isokinetic (fulfills criteria)
- 3) Satisfactory shoulder stability
- 4) No pain or tenderness

#### Exercises:

- Gradually progress sport activities to unrestrictive participation
- Continue stretching and strengthening program