

Therapist _____

Phone

Type Two Rotator Cuff Repair Arthroscopic Assisted with SAD Medium to Large Tears (Greater than 1 cm, Less than 5 cm)

I. Phase I - Immediate Post-Surgical Phase (Day 1-10)

<u>Goals</u>: Maintain Integrity of the Repair Gradually Increase Passive Range of Motion Diminish Pain and Inflammation Prevent Muscular Inhibition

Day One to Six:

- Sling or Slight Abduction Brace **Physician Decision
- Pendulum Exercises 4-8x daily (flexion, circles)
- Active Assisted ROM Exercise (L-Bar)
 - ER/IR in Scapular Plane
 - Passive ROM
 - Flexion to tolerance
 - ER/IR in Scapular Plane
- Elbow/Hand Gripping & ROM Exercises
 - Submaximal Painfree Isometrics
 - Flexion
 - Abduction
 - External Rotation
 - Internal Rotation
 - Elbow Flexors
- Cryotherapy for Pain and Inflammation
 - Ice 15-20 minutes every hour
- Sleeping
 - Sleep in Sling or Brace

Day Seven to Ten:

- Continue Pendulum Exercises
- Progress Passive ROM to Tolerance
 - Flexion to at least 105 degrees
 - ER in Scapular Plane to 35-45 degrees
 - IR in Scapular Plane to 35-45 degrees
- Active Assisted ROM Exercises (L-bar)
 - ER/IR in Scapular Plane





- Flexion to Tolerance*
- *Therapist Provides Assistance by Supporting Arm
- Continue Elbow/Hand ROM & Gripping Exercises
- Continue Isometrics
 - Flexion with Bent Elbow
 - Extension with Bent Elbow
 - Abduction with Bent Elbow
 - ER/IR with Arm in Scapular Plane
 - Elbow Flexion
- Continue Use of Ice for Pain Control
 - Use Ice at least 6-7 times daily
- Sleeping
 - Continue Sleeping in Brace until Physician Instructs

Precautions:

- 1. No Lifting of Objects
- 2. No Excessive Shoulder Extension
- 3. No Excessive Stretching or Sudden Movements
- 4. No Supporting of Body Weight by Hands
- 5. Keep Incision Clean & Dry

II. Phase II - Protection Phase (Day 11 – Week 6)

Goals: Allow Healing of Soft Tissue

Do Not Overstress Healing Tissue Gradually Restore Full Passive ROM (Week 4-5) Re-Establish Dynamic Shoulder Stability Decrease Pain & Inflammation

<u>Day 11 – 14:</u>

- Passive Range of Motion to Tolerance
 - Flexion 0-125/145 degrees
 - ER at 90 degrees abduction to at least 45 degrees
 - IR at 90 degrees abduction to at least 45 degrees
- Active Assisted ROM to Tolerance
 - Flexion
 - ER/IR in Scapular Plane
 - ER/IR at 90 degrees Abduction
- Dynamic Stabilization Drills
 - Rhythmic Stabilization Drills
 - ER/IR in Scapular Plane

AOKC.NET



- Flexion/Extension at 100 degrees Flexion
- Continue All Isometric Contractions
- Continue Use of Cryotherapy as needed
- Continue All Precautions

Week 3 - 4:

- Patient should exhibit full passive range of motion
- Continue all exercises listed above
- Initiate ER/IR strengthening using exercise tubing at 0 degrees of abduction
- Initiate Manual Resistance ER Supine in Scapular Plane
- Initiate Prone Rowing to Neutral arm Position
- Initiate Isotonic Elbow Flexion
- Continue use of ice as needed
- May use heat prior to ROM exercises
- May use pool for light ROM exercises

<u>Week 5 – 6:</u>

- May use heat prior to exercises
- Continue AAROM and Stretching exercises
- Initiate Active ROM Exercises
 - Shoulder Flexion Scapular Plane
 - Shoulder Abduction
- Progress Isotonic Strengthening Exercise Program
 - ER Tubing
 - Sidelying ER
 - Prone Rowing
 - Prone Horizontal Abduction
 - Biceps Curls

Precautions:

- 1. No Heavy Lifting of Objects
- 2. No excessive behind the back movements
- 3. No Supporting of Body Weight by Hands & Arms

AOKC.NET

4. No Sudden Jerking Motions

III. Phase III – Intermediate Phase (Week 7-14)

<u>Goals</u>: Full Active ROM (Week 8-10) Full Passive ROM Dynamic Shoulder Stability Gradual Restoration of Shoulder Strength & Power Gradual Return to Functional Activities



Week 7:

- Continue Stretching & PROM (as needed to maintain full ROM)
- Continue Dynamic Stabilization Drills
- Progress Strengthening Program
 - ER/IR Tubing
 - ER Sidelying
 - Lateral Raises*
 - Full Can in Scapular Plane*
 - Prone Rowing
 - Prone Horizontal Abduction
 - Prone Extension
 - Elbow Flexion
 - Elbow Extension

*Patient must be able to elevate arm without shoulder or scapular hiking before initiating isotonics; if unable, continue glenohumeral dynamic stabilization exercises.

Week 8:

- Continue all exercise listed above
- If physician permits, may initiate Light functional activities

Week 14:

- Continue all exercise listed above
- Progress to Independent Home Exercise Program (Fundamental Shoulder Exercises)

IV. <u>Phase IV</u> – <u>Advanced Strengthening Phase (Week 15 - 22)</u>

<u>Goals</u>: Maintain Full Non-Painful ROM Enhance Functional Use of UE Improve Muscular Strengthen & Power Gradual Return to Functional Activities

Week 15:

- Continue ROM & Stretching to maintain full ROM
- Self Capsular Stretches
- Progress Shoulder Strengthening Exercises
 - Fundamental Shoulder Exercises
- Initiate Interval Golf Program (if appropriate)

<u>Week 20:</u>

- Continue all exercises listed above
- Progress Golf Program to playing golf (if appropriate)
- Initiate Interval Tennis Program (if appropriate)
- May Initiate Swimming

AOKC.NET



V. <u>Phase V</u> – <u>Return to Activity Phase (Week 23 - 30)</u>

<u>Goals</u>: Gradual Return to Strenuous Work Activities Gradual Return to Recreational Sport Activities

Week 23:

- Continue Fundamental Shoulder Exercise Program (at least 4 times weekly)
- Continue Stretching, if motion is tight
- Continue Progression to Sport Participation

AOKC.NET